

4F13: Machine Learning

A Belief Propagation Demo

Zoubin Ghahramani

zoubin@eng.cam.ac.uk


**Department of Engineering
University of Cambridge**

Michaelmas, 2006

<http://learning.eng.cam.ac.uk/zoubin/ml06/>


1. Model Structure

Fluffy = pet cat
Moby = pet fish




2. Model Parameters

Fluffy = pet cat
Moby = pet fish



3. Propagating Evidence




1. Observe "Moby is dead", i.e. $x_3 = 1$

2. Send $\lambda_{x_3}(x_1) \equiv p(e_{x_1 \rightarrow x_3}^- | x_1) = \begin{bmatrix} 0.1 \\ 1.0 \end{bmatrix}$ message $x_3 \rightarrow x_1$

3. $BEL(x_1 | x_3 = 1) = \frac{1}{Z} \begin{bmatrix} 0.99 \\ 0.01 \end{bmatrix} \odot \begin{bmatrix} 0.1 \\ 1.0 \end{bmatrix} = \begin{bmatrix} 0.91 \\ 0.09 \end{bmatrix}$

4. Propagating Evidence




4. Send $\pi_{x_4}(x_1) \equiv p(x_1 | e_{x_1 \rightarrow x_4}^+) = \begin{bmatrix} 0.91 \\ 0.09 \end{bmatrix}$

5. Send $\pi_{x_4}(x_2) \equiv p(x_2 | e_{x_2 \rightarrow x_4}^+) = p(x_2) = \begin{bmatrix} 0.99 \\ 0.01 \end{bmatrix}$ from $x_2 \rightarrow x_4$.

6. Compute $\pi(x_4) \equiv p(x_4 | e_{x_4}^+) = \sum_{x_1, x_2} p(x_4 | x_1, x_2) \pi_{x_4}(x_1) \pi_{x_4}(x_2) = \begin{bmatrix} 0.18 \\ 0.82 \end{bmatrix}$

7. $BEL(x_4 | x_3 = 1) = \begin{bmatrix} 0.18 \\ 0.82 \end{bmatrix}$, whereas before observing $x_3 = 1$, $BEL(x_4) = \begin{bmatrix} 0.1 \\ 0.9 \end{bmatrix}$.

5. Propagating Evidence



8. Observe “Fluffy’s Food Bowl is Full” $x_6 = 1$!

9. Send $\lambda_{x_6}(x_4) = \begin{bmatrix} 0.9 \\ 0.1 \end{bmatrix}$ message $x_6 \rightarrow x_4$

10. $BEL(x_4|x_3 = 1, x_6 = 1) = \frac{1}{Z} \begin{bmatrix} 0.18 \\ 0.82 \end{bmatrix} \odot \begin{bmatrix} 0.9 \\ 0.1 \end{bmatrix} = \begin{bmatrix} 0.66 \\ 0.34 \end{bmatrix}$

11. Send $\lambda_{x_4}(x_1) = \sum_{x_4} \lambda_{x_6}(x_4) \sum_{x_2} p(x_4|x_1, x_2) \pi_{x_4}(x_2) = \begin{bmatrix} 0.19 \\ 0.82 \end{bmatrix}$

12. $BEL(x_1|x_3 = 1, x_6 = 1) = \frac{1}{Z} \begin{bmatrix} 0.99 \\ 0.01 \end{bmatrix} \odot \begin{bmatrix} 0.1 \\ 1.0 \end{bmatrix} \odot \begin{bmatrix} 0.19 \\ 0.82 \end{bmatrix} = \begin{bmatrix} 0.70 \\ 0.30 \end{bmatrix} \Rightarrow$ Fluffy still innocent!